Fluid Phase Equilibria, 81 (1992) 357–358
Elsevier Science Publishers B.V., Amsterdam

Contents Volume 81 (1992)

The predictive capability of the UNIFAC equation for the calculation of liquid–liquid equilibria with chemical reactions in tributyl phosphate extraction systems for nitric acid and uranyl nitrate
J. Chen, Z. C. Li, J. F. Lu, and Y. G. Li (Beijing, China) 1–15

Application of the modified Kumar–Starling equation of state to predict the phase behavior of mixtures in the near-critical region
J.-Z. Chu, Y.-X. Zuo, and T.-M. Guo (Beijing, China)..... 17–38

Critical point calculations for oil reservoir fluid systems using the SPHCT equation of state
F. García-Sánchez, J. L. Ruiz-Cortina, C. Lira-Galeana and L. Ponce-Ramirez (Mexico City, Mexico)..... 39–84

Standard states, reference states and finite-concentration effects in near-critical mixtures with applications to aqueous solutions
J.M.H. Levelt Sengers (Gaithersburg, MD, USA), A. H. Harvey (Fullerton, CA, USA), R. Crovetto (University Park, PA, USA) and J. S. Gallagher (Gaithersburg, MD, USA)..... 85–107

Extension of the group contribution equation of state for the calculation of gas solubilities
S.R. Wolff, R. P. Danner (University Park, PA, USA) and Aa. Fredenslund (Lyngby, Denmark) 109–127

Estimation of interfacial tensions of ternary liquid–liquid systems by UNIQUAC method
B. Li and J. Fu (Beijing, China) 129–152

A new method to construct equations of state and some examples
H.-Y. Lee (Shanghai, China)..... 153–164

Vapour pressures of butan-1-ol with *n*-hexadecane between 293.18 and 323.18 K. Description of butan-1-ol + *n*-alkane systems by ERAS model
I. Baños (Logroño, Spain), F. Sánchez, P. Pérez, J. Valero and M. Gracia (Zaragoza, Spain) 165–174

High pressure vapor liquid equilibria at 293 K for systems containing nitrogen, methane and carbon dioxide
N. Xu, J. Dong, Y. Wang and J. Shi (Nanjing, China) 175–186

An examination of the vapor–liquid equilibrium in the system propane–hydrogen sulfide
J.J. Carroll and A.E. Mather (Edmonton, Alb., Canada)..... 187–204

Vapour–liquid and liquid–liquid equilibria for the propane–furfural and propylene–furfural systems
K. Noda, H. Watanabe and S. Iwamoto (Hamamatsu, Japan) 205–215

Phase equilibria in very dilute mixtures of water and unsaturated chlorinated hydrocarbons and of water and benzene
M.R. Cooling, B. Khalfoui and D.M.T. Newsham (Manchester, UK) 217–229

Excess molar enthalpies of binary mixtures ethylbenzene + propylbenzene + butylbenzene and + hexylbenzene at 298.15, 308.15 and 318.15 K
D.V.S. Jain and N.S. Dhar (Chandigarh, India)..... 231–239

Solubilities of hydrogen cyanide and sulfur dioxide in water at temperatures from 293.15 to 413.15 K and pressures up to 2.5 MPa
B. Rumpf and G. Maurer (Kaiserslautern, Germany) 241–260
Vapour–liquid equilibria. VI. The ternary system acetone–hexane–cyclohexane at 313.15 K
M. Góral, P. Oracz and S. Warycha (Warsaw, Poland) 261–272
Excess molar heat capacities of binary mixtures of ethylbenzene with xylenes from 303.65 to 321.65 K
D.V.S. Jain, R. Chadha and S.K. Sehgal (Chandigarh, India)..... 273–283
Thermodynamic properties of $CHF_2–O–CHF_2$, bis(difluoromethyl) ether
D.R. Defibaugh, K.A. Gillis, M.R. Moldover, G. Morrison and J. W. Schmidt (Gaithersburg, MD, USA) 285–305
Phase equilibrium study for the separation and fractionation of fatty oil components using supercritical carbon dioxide
R. Bharath, H. Inomata, T. Adshiri and K. Arai (Sendai, Japan) . 307–320
Multiphase behavior of binary and ternary systems of heavy aromatic hydrocarbons with supercritical carbon dioxide. Part I. Experimental results
S.T. Chung and K.S. Shing (Los Angeles, CA, USA)..... 321–341

Letters to the Editor

Comments concerning "On the isobaric vapour–liquid equilibrium of the cyclohexanol–phenylmethanol binary at 101.325 ± 0.067 kPa"
W.E. Acree, Jr. (Denton, TX, USA)..... 343–345
On the consistency of SEMAC equations
A.A. Sayar (Istanbul, Turkey) 347–348

Corrigenda 349

Announcement 351–353

Author Index 355

Contents of Volume 81 (1992)..... 357–358